

PRESCRIBED USE

- To decrease restlessness in patients that are overactive, cannot concentrate or are easily distracted.
- Often used in combination with social, educational & psychological therapy

WHY IS IT A CONTROLLED SUBSTANCE/ YOU NEED A PRESCRIPTION FOR IT?

- Can cause psychosis, delirium, irregular heartbeat, panic and heart failure
- Therefore dangerous if used improperly and without physician oversight.

OTHER INFO

- Stimulant
- Treats ADHD & Narcolepsy
- Addictive

EXTREMELY DANGEROUS TO TAKE IF YOU HAVE THESE ILLNESSES:

- Cardiac Disease
- Hypertension
- Seizure Disorders
- Substance Abuse
- Liver disease
- Glaucoma

WARNING

Should NOT be taken with antidepressants! (i.e. Prozac and Cymbalta)

ADDERALL

STREET NAMES

- Pep Pills
- Addy
- Uppers
- Black Beauties

SERIOUS SIDE EFFECTS

- Blurred vision
- Allergic reaction
- Chest pain, trouble breathing, fainting
- Extreme energy, unusual behavior, confusion, agitation
- Fast, pounding uneven heartbeat
- Numb, cold, pale or painful fingers or toes
- Delusions & Seizures

LESS SERIOUS SIDE EFFECTS

- Dry mouth, diarrhea, stomach pain, nausea
- Headache, dizziness
- Loss of appetite, weight-loss
- Trouble sleeping
- Change in libido
- Hair loss

